





Electron emission modeling in ionisation gauges for magnetic confinement fusion

IPP

A. Glock*, M. Griener, C. Drobny,
G. Birkenmeier*, H. Meister, G. Schlisio

Max Planck Institute for Plasma Physics, Garching & Greifswald, Germany

*Technische Universität München, TUM School of Natural Sciences., Physics Dep.,
Garching, Germany



This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.




Federal Ministry
of Research, Technology
and Space

This work received funding from the German
Ministry of Research, Technology and Space
(BMFT) within the funding program Fusion
2040 (reference no. 13F1020).



Magnetic confinement fusion research at IPP


Wendelstein 7-X Stellarator (Greifswald)



- Operating since 2015
- Plasma volume $\sim 30\text{m}^3$
- Magnetic field 3T (on axis)



ASDEX Upgrade Tokamak (Garching)



- Operating since 1991
- Plasma volume $\sim 13\text{m}^3$
- Magnetic field 2.5T (on axis)





In vessel pressure measurements

In vessel pressure for **real-time plasma control**.

Conditions close to the plasma, behind first wall:

- **Pressure range** from 10^{-4} – 30Pa
- **Time resolution** $< 1\text{ms}$
- **Magnetic field** 1 – 8T (tested range)
- **Neutron flux / radiation** 10^2 – 10^5Gy/h (ITER)
- **Temperature** up to 700K (ITER)

→ **Not possible with Pirani, Baratron and cold cathode gauges**


→ still used for machine safety and calibration



**„ASDEX“-type
pressure gauge**




Functionality:







ASDEX gauge – hardware



hot cathode ionization gauge





ASDEX gauge - potentials



Based on: Haas, G. & Bosch, H.-S. *Vacuum* 51, 39–46 (1998)




ASDEX gauge - output



Based on: Haas, G. & Bosch, H.-S. *Vacuum* 51, 39–46 (1998)

Modeling:



Overview




Design optimization and “synthetic diagnostic”:

- Performance prediction and improvement
- Parameter sweeps and automation
- Extrapolation beyond lab capabilities

Characteristic plasma physics:

- Access to “invisible” quantities
- Isolated effects and Laboratory verification


Focusing on collisionless electrons (for now)





Thermionic emission & electron dynamics


Input:



- No gas background = **no ions**
- Sinusoidal variation in filament heating

Experimental observation:

A. C. Castillo et al., *Fusion Engineering and Design*, Volume 181 (2022)




- Grid current follows heating power
- **Reproducible, hysteretic jumps**

→ State transitions/perturbations in the electron population




“Electromagnetostatic” Particle-In-Cell method



Used codes



Geometry generation:



Gmsh [1]

- Generating complex geometries in 3D


PIC simulation:



[2]

- Massively parallelized with openMP/MPI
- Scalable on CPU and GPU clusters

Postprocessing:



- Matplotlib for standard 2D plots
- Paraview for 3D and particle visualization

[1] C. Geuzaine and J.-F. Remacle, **Gmsh**: a three-dimensional finite element mesh generator with built-in pre- and post-processing facilities, International Journal for Numerical Methods in Engineering, Volume 79 (2009)


[2] J.-L. Vay et al., **WarpX**: An advanced Particle-In-Cell code. [DOI:10.5281/zenodo.4571577](https://doi.org/10.5281/zenodo.4571577), <https://blast-warpX.github.io> (2018)

This research used the open-source [particle-in-cell code WarpX](#). Primary WarpX contributors are with LBNL, LLNL, CEA-LIDYL, SLAC, DESY, CERN, Helion Energy, TAE Technologies, and Realta Fusion. We acknowledge all WarpX contributors.




The electron model

1. General constraints




$\Delta t, \Delta x$


2. Emitter types




3. Source flux



4. Source distributions





5. Boundaries





1. General constraints



1. Debye and geometric:

$$\Delta x \approx \mathcal{O}(10^{-4}\text{m})$$


2. Maximum particle velocity:

$$v_{\text{max}} \approx \sqrt{\frac{2 q_e U_{\text{max}}}{m_e}} + 4 v_{\text{th}}$$


$$\Delta t \approx \mathcal{O}(10^{-12}\text{s})$$




2. Emitter types




Spiral design



Meander design



LaB₆



ZrC

Actively heated emitters

Passively heated emitters


* Filament diameters not to scale



3. Source flux

Using the **Richardson-Dushman** law:


$$J_s = A_G T^2 \exp\left(-\frac{W_f}{k_B T}\right) C(T), \quad \sqrt{\frac{q_e^3 E}{4\pi\epsilon_0}} \approx \text{const.}$$



And the most probable **thermal velocity**:


$$v_{th} = \sqrt{\frac{k_B T}{m_e}}$$

Cutoff at $\pm 4v_{th}$ (99.38% of velocity dist.)





4. Source distribution




- Spatial non-uniform source flux

- Partial surfaces



5. Boundary conditions



Electrostatics:

- **Fixed potentials** as Dirichlet conditions:

$$\phi|_{\partial\Omega} = U$$

Particles:

- **Absorbing boundaries** (perfect conductor)


$$I|_{\partial\Omega} = \frac{dQ}{dt}|_{\partial\Omega}$$

Mirror plane:

- **Neumann potential**
- **Inversion of normal particle velocity**





First results – magnetic field influence





no magnetic field

perfectly aligned field of 1T





First results – electron trapping and current oscillations





Conclusion

Space-charge formation:

- The gauge geometry traps electrons, even for relatively weak background fields at the ends of the oscillation volume.
- Electron densities continue to grow for more **>100 electron oscillation cycles**, reaching higher peak values than initially expected.

“Steady-state” oscillations & current instability:

- When is the true steady state reached - **μs or ms timescales?** How does this depend on the **B-field magnitude and gauge parameters/emitters?**
 - What is the driving mechanism behind the observed oscillations?
- **Further analysis and follow-up studies of source transients are planned.**

For more information and/or questions: alexander.glock@ipp.mpg.de